arXiv:0912.4246v1 [math.AP] 21 Dec 2009

PERTURBATION METHOD FOR PARTICLE-LIKE SOLUTIONS OF THE EINSTEIN-DIRAC-MAXWELL EQUATIONS

SIMONA ROTA NODARI

Abstract. The aim of this Note is to prove by a perturbation method the existence of solutions of the coupled Einstein-Dirac-Maxwell equations for a static, spherically symmetric system of two fermions in a singlet spinor state and with the electromagnetic coupling constant $\left(\frac{e}{m}\right)^2 < 1$. We show that the nondegenerate solution of Choquard's equation generates a branch of solutions of the Einstein-Dirac-Maxwell equations.

Une méthode de perturbation pour les solutions localisées des équations d'Einstein-Dirac-Maxwell.

Résumé. Le but de cette Note est de démontrer par une méthode de perturbation l'existence de solutions des équations d'Einstein-Dirac-Maxwell pour un système statique, à symétrie sphérique de deux fermions dans un état de singulet et avec une constante de couplage électromagnétique $\left(\frac{e}{m}\right)^2 < 1$. On montre que la solution non dégénérée de l'équation de Choquard génère une branche de solutions des équations d'Einstein-Dirac-Maxwell.

Version française abrégée

Dans un papier récent [5], par une méthode de perturbation, on a montré de manière rigoureuse l'existence de solutions des équations d'Einstein-Dirac pour un système statique, à symétrie sphérique de deux fermions dans un état de singulet. Dans cette Note, on généralise ce résultat aux équations d'Einstein-Dirac-Maxwell et on montre, dans le cas particulier d'un couplage électromagnétique faible, l'existence des solutions obtenues numériquement par F. Finster, J. Smoller et ST. Yau dans [7].

Plus précisément, en utilisant l'idée introduite par Ounaies pour une classe d'équations de Dirac non linéaires (voir [1]) et adaptée dans [5] aux équations d'Einstein-Dirac, on obtient le théorème suivant.

Théorème 0.1. *Soient e, m, ω tels que $e^2 - m^2 < 0$, $0 < \omega < m$ et supposons $m - \omega$ assez petit ; alors il existe une solution non triviale de (1-5).*

Dans cette Note, on décrit la méthode utilisée pour démontrer ce théorème. Premièrement, par un changement d'échelle, on transforme les équations d'Einstein-Dirac-Maxwell (6-9) en un système perturbé qui s'écrit sous la forme (11). On choisit $\varepsilon = m - \omega$ comme paramètre de perturbation.

Deuxièment, on remarque que, pour $\varepsilon = 0$ et $\left(\frac{e}{m}\right)^2 < 1$, ce système est équivalent au système (12) où l'équation pour la variable φ est l'équation de Choquard. Il est bien connu que l'équation de Choquard a une solution radiale positive. De plus, dans l'espace des fonctions radiales, cette solution est non dégénérée, dans le sens où le

---

Date: December 18, 2009.

noyau de la linéarisation de l'équation contient seulement la fonction identiquement nulle. On appelle ϕ_0 la solution du système (12).

Ensuite, on observe que le système perturbé s'écrit sous la forme $D(\varepsilon,\varphi,\chi,\tau,\zeta)=0$ avec D un opérateur non linéaire de classe $\mathcal{C}^1$, pour un bon choix d'espaces fonctionnels. On prouve que cet opérateur satisfait les hypothèses du théorème des fonctions implicites. En particulier, on montre que la linéarisation de l'opérateur D par rapport à $(\varphi,\chi,\tau,\zeta)$ en $(0,\phi_0)$, $D_{\varphi,\chi,\tau,\zeta}(0,\phi_0)$, est une injection, grâce à la non-dégénérescence de la solution de l'équation de Choquard, et s'écrit comme somme d'un isomorphisme et d'un opérateur compact ; donc $D_{\varphi,\chi,\tau,\zeta}(0,\phi_0)$ est un isomorphisme. En appliquant le théorème des fonctions implicites, on déduit que, pour ε assez petit et $e^2-m^2<0$, le système (11) a une solution.

En conclusion, pour $e^2-m^2<0$, $0<\omega<m$ et $m-\omega$ assez petit, les équations d'Einstein-Dirac-Maxwell possèdent une solution non triviale.

1. Introduction

In a recent paper [5], using a perturbation method, we proved rigorously the existence of solutions of the coupled Einstein-Dirac equations for a static, spherically symmetric system of two fermions in a singlet spinor state. In this Note, we extend our result to the Einstein-Dirac-Maxwell equations and we prove, in the particular case of a weak electromagnetic coupling, the existence of the solutions obtained numerically by F. Finster, J. Smoller and ST. Yau in [7].

The general Einstein-Dirac-Maxwell equations for a system of n Dirac particles take the form

$$(G-m)\psi_a=0,\quad R^i_j-\tfrac{1}{2}R\delta^i_j=-8\pi T^i_j,\quad \nabla_k F^{jk}=4\pi e\sum_{a=1}^{n}\overline{\psi_a}G^j\psi_a$$

where G^j are the Dirac matrices, G denote the Dirac operator, ψ_a are the wave functions of fermions of mass m and charge e, F_{jk} is the electromagnetic field tensor and, finally, T^i_j is the sum of the energy-momentum tensor of the Dirac particle and the Maxwell stress-energy tensor.

In [7], the metric, in polar coordinates (t,r,ϑ,φ), is given by

$$ds^2=\frac{1}{T^2}\,dt^2-\frac{1}{A}\,dr^2-r^2\,d\vartheta^2-r^2\sin^2\vartheta\,d\varphi^2$$

with $A=A(r)$, $T=T(r)$ positive functions; moreover, using the ansatz from [6], Finster, Smoller and Yau describe the Dirac spinors with two real radial functions $\Phi_1(r)$, $\Phi_2(r)$ and they assume that the electromagnetic potential has the form $\mathcal{A}=(-V,0)$, with V the Coulomb potential.

In this case the Einstein-Dirac-Maxwell equations can be written as

$$\begin{aligned}
\sqrt{A}\Phi_1' &= \frac{1}{r}\Phi_1 - ((\omega - eV)T + m)\Phi_2 && (1)\\
\sqrt{A}\Phi_2' &= ((\omega - eV)T - m)\Phi_1 - \frac{1}{r}\Phi_2 && (2)\\
rA' &= 1 - A - 16\pi(\omega - eV)T^2\left(\Phi_1^2 + \Phi_2^2\right) - r^2AT^2\left(V'\right)^2 && (3)\\
2rA\frac{T'}{T} &= A - 1 - 16\pi(\omega - eV)T^2\left(\Phi_1^2 + \Phi_2^2\right) + 32\pi\frac{1}{r}T\Phi_1\Phi_2 \\
&\quad +16\pi mT\left(\Phi_1^2 - \Phi_2^2\right) + r^2AT^2\left(V'\right)^2 && (4)\\
r^2AV'' &= -8\pi e\left(\Phi_1^2 + \Phi_2^2\right) - \left(2rA + r^2A\frac{T'}{T} + \frac{r^2}{2}A'\right)V' && (5)
\end{aligned}$$

with the normalization condition $\int_0^\infty |\Phi|^2 \frac{T}{\sqrt{A}}\,dr = \frac{1}{4\pi}$.
In order that the metric be asymptotically Minkowskian and the solutions have finite (ADM) mass, Finster, Smoller and Yau assume

$$\lim_{r\to\infty} T(r) = 1$$

and

$$\lim_{r\to\infty} \frac{r}{2}(1 - A(r)) < \infty.$$

Finally, they also require that the electromagnetic potential vanishes at infinity.

In this Note, using the idea introduced by Ounaies for a class of nonlinear Dirac equations (see [1]) and adapted in [5] to the Einstein-Dirac equations, we obtain the following result.

Theorem 1.1. *Given e, m, ω such that $e^2 - m^2 < 0$, $0 < \omega < m$ and $m - \omega$ is sufficiently small, there exists a non trivial solution of (1-5).*

2. PERTURBATION METHOD FOR THE EINSTEIN-DIRAC-MAXWELL EQUATIONS

First of all, we observe that, writing $T(r) = 1 + t(r)$ and integrating the equation (5), the Einstein-Dirac-Maxwell equations become

$$\begin{aligned}
\sqrt{A}\Phi_1' &= \frac{1}{r}\Phi_1 - ((\omega - eV)(1+t) + m)\Phi_2 && (6)\\
\sqrt{A}\Phi_2' &= ((\omega - eV)(1+t) - m)\Phi_1 - \frac{1}{r}\Phi_2 && (7)\\
2rAt' &= (A-1)(1+t) - 16\pi(\omega - eV)(1+t)^3\left(\Phi_1^2 + \Phi_2^2\right) \\
&\quad +32\pi\frac{1}{r}(1+t)^2\Phi_1\Phi_2 + 16\pi m(1+t)^2\left(\Phi_1^2 - \Phi_2^2\right) \\
&\quad +r^2A(1+t)^3\left(V'\right)^2 && (8)\\
\sqrt{A}(1+t)V' &= -\frac{8\pi e}{r^2}\int_0^r \left(\Phi_1^2 + \Phi_2^2\right)\frac{(1+t)}{\sqrt{A}}\,ds. && (9)
\end{aligned}$$

where $A(r) = 1 + a(r)$ and

$$\begin{aligned}
a(r) &= -\frac{1}{r}\exp\left(-F(r)\right)\int_0^r \Big[16\pi(\omega - eV)(1+t)^2\left(\Phi_1^2 + \Phi_2^2\right) \\
&\quad +s^2(1+t)^2\left(V'\right)^2\Big]\exp\left(F(s)\right)ds && (10)
\end{aligned}$$

with $F(r)=\int_0^r s(1+t)^2\left(V'\right)^2\,ds$.

After that, we introduce the new variable $(\varphi,\chi,\tau,\zeta)$ such that

$$\Phi_1(r)=\varepsilon^{1/2}\varphi(\varepsilon^{1/2}r),\quad \Phi_2(r)=\varepsilon\chi(\varepsilon^{1/2}r),\quad t(r)=\varepsilon\tau(\varepsilon^{1/2}r),\quad V(r)=\varepsilon\zeta(\varepsilon^{1/2}r)$$

where Φ_1,Φ_2,t,V satisfy (6-9) and $\varepsilon=m-\omega$. Using the explicit expression of $a(r)$, given in (10), we write

$$a(\Phi_1,\Phi_2,t,V)=\varepsilon\alpha(\varepsilon,\varphi,\chi,\tau,\zeta)$$

with $\alpha(0,\varphi,\chi,\tau,\zeta)=-\frac{16\pi m}{r}\int_0^r\varphi^2\,ds$. It is now clear that if Φ_1,Φ_2,t,V satisfy (6-9), then φ,χ,τ,ζ satisfy the system

$$\begin{cases}(1+\varepsilon\alpha(\varepsilon,\varphi,\chi,\tau,\zeta))^{1/2}\frac{d}{dr}\varphi-\frac{1}{r}\varphi+2m\chi+K_1\left(\varepsilon,\varphi,\chi,\tau,\zeta\right)=0\\ (1+\varepsilon\alpha(\varepsilon,\varphi,\chi,\tau,\zeta))^{1/2}\frac{d}{dr}\chi+\frac{1}{r}\chi+\varphi-m\varphi\tau+e\varphi\zeta+K_2\left(\varepsilon,\varphi,\chi,\tau,\zeta\right)=0\\ (1+\varepsilon\alpha(\varepsilon,\varphi,\chi,\tau,\zeta))\frac{d}{dr}\tau-\frac{\alpha(\varepsilon,\varphi,\chi,\tau,\zeta)}{2r}+K_3\left(\varepsilon,\varphi,\chi,\tau,\zeta\right)=0\\ (1+\varepsilon\alpha(\varepsilon,\varphi,\chi,\tau,\zeta))^{1/2}(1+\varepsilon\tau)\frac{d}{dr}\zeta+\frac{8\pi e}{r^2}\int_0^r\varphi^2\,ds+K_4\left(\varepsilon,\varphi,\chi,\tau,\zeta\right)=0\end{cases}\tag{11}$$

where $K_1\left(0,\varphi,\chi,\tau,\zeta\right)=K_2\left(0,\varphi,\chi,\tau,\zeta\right)=K_3\left(0,\varphi,\chi,\tau,\zeta\right)=K_4\left(0,\varphi,\chi,\tau,\zeta\right)=0$.

Then, for $\varepsilon=0$, (11) becomes

$$\begin{cases}-\frac{d^2}{dr^2}\varphi+2m\varphi+16\pi(e^2-m^2)m\left(\int_0^\infty\frac{\varphi^2}{\max(r,s)}\,ds\right)\varphi=0\\ \chi(r)=\frac{1}{2m}\left(\frac{1}{r}\varphi-\frac{d}{dr}\varphi\right)\\ \tau(r)=8\pi m\int_0^\infty\frac{\varphi^2}{\max(r,s)}\,ds\\ \zeta(r)=8\pi e\int_0^\infty\frac{\varphi^2}{\max(r,s)}\,ds\end{cases}\tag{12}$$

We remark that if $e^2-m^2<0$ the first equation of the system (12) is the Choquard equation

$$-\triangle u+2mu-4(m^2-e^2)m\left(\int_{\mathbb{R}^3}\frac{|u(y)|^2}{|x-y|}\,dy\right)u=0\quad\text{in } H^1\left(\mathbb{R}^3\right)\tag{13}$$

with $u(x)=\frac{\varphi(|x|)}{|x|}$. It is well known that Choquard's equation (13) has a unique radial, positive solution u_0 with $\int|u_0|^2=N$ for some $N>0$ given. Furthermore, u_0 is infinitely differentiable, goes to zero at infinity and is a radial nondegenerate solution; by this we mean that the linearization of (13) around u_0 has a trivial nullspace in $L^2_r(\mathbb{R}^3)$ (see [3], [4], [2] for more details).

Let $\phi_0=(\varphi_0,\chi_0,\tau_0,\zeta_0)$ be the ground state solution of (12).

The main idea is that the solutions of (11) are the zeros of a $\mathcal{C}^1$ operator $D:\mathbb{R}\times X_\varphi\times X_\chi\times X_\tau\times X_\zeta\to Y_\varphi\times Y_\chi\times Y_\tau\times Y_\zeta$. So, to obtain a solution of (11) from

ϕ_0, we define the operators

$$\begin{aligned}
L_1(\varepsilon,\varphi,\chi,\tau,\zeta) &= (1+\varepsilon\alpha(\varepsilon,\varphi,\chi,\tau,\zeta))^{1/2}\frac{1}{r}\frac{d}{dr}\varphi-\frac{\varphi}{r^2}+2m\frac{\chi}{r}\\
&\quad+\frac{1}{r}K_1\left(\varepsilon,\varphi,\chi,\tau,\zeta\right)\\
L_2(\varepsilon,\varphi,\chi,\tau,\zeta) &= (1+\varepsilon\alpha(\varepsilon,\varphi,\chi,\tau,\zeta))^{1/2}\frac{1}{r}\frac{d}{dr}\chi+\frac{\chi}{r^2}+\frac{\varphi}{r}-m\frac{\varphi}{r}\tau+e\frac{\varphi}{r}\zeta\\
&\quad+\frac{1}{r}K_2\left(\varepsilon,\varphi,\chi,\tau,\zeta\right)\\
L_3(\varepsilon,\varphi,\chi,\tau,\zeta) &= (1+\varepsilon\alpha(\varepsilon,\varphi,\chi,\tau,\zeta))\frac{d}{dr}\tau-\frac{\alpha(\varepsilon,\varphi,\chi,\tau,\zeta)}{2r}+K_3\left(\varepsilon,\varphi,\chi,\tau,\zeta\right)\\
L_4(\varepsilon,\varphi,\chi,\tau,\zeta) &= (1+\varepsilon\alpha(\varepsilon,\varphi,\chi,\tau,\zeta))^{1/2}(1+\varepsilon\tau)\frac{d}{dr}\zeta+\frac{8\pi e}{r^2}\int_0^r\varphi^2\,ds\\
&\quad+K_4\left(\varepsilon,\varphi,\chi,\tau,\zeta\right)
\end{aligned}$$

and

$$D(\varepsilon,\varphi,\chi,\tau,\zeta)=\left(L_1(\varepsilon,\varphi,\chi,\tau,\zeta),L_2(\varepsilon,\varphi,\chi,\tau,\zeta),L_3(\varepsilon,\varphi,\chi,\tau,\zeta),L_4(\varepsilon,\varphi,\chi,\tau,\zeta)\right),$$

with X_φ, X_χ, X_τ, $Y_\varphi, Y_\chi, Y_\tau$ defined as in [5] and

$$\begin{aligned}
X_\zeta &= \left\{\zeta:(0,\infty)\to\mathbb{R}\,\middle|\,\lim_{r\to\infty}\zeta(r)=0,\ \frac{d}{dr}\zeta\in L^1((0,\infty),dr)\cap L^2((0,\infty),rdr)\right\}\\
Y_\zeta &= L^1((0,\infty),dr)\cap L^2((0,\infty),rdr)
\end{aligned}$$

with their natural norms.

Next, we linearize the operator D on $(\varphi,\chi,\tau,\zeta)$ around $(0,\phi_0)$:

$$\begin{aligned}
D_{\varphi,\chi,\tau,\zeta}(0,\phi_0)(h,k,l,z) &= \begin{pmatrix}\frac{1}{r}\frac{d}{dr}h-\frac{h}{r^2}+2m\frac{k}{r}\\ \frac{1}{r}\frac{d}{dr}k+\frac{k}{r^2}+\frac{h}{r}-m\frac{\varphi_0}{r}l+e\frac{\varphi_0}{r}z\\ \frac{d}{dr}l\\ \frac{d}{dr}z\end{pmatrix}\\
&\quad+\begin{pmatrix}0\\ -m\frac{h}{r}\tau_0+e\frac{h}{r}\zeta_0\\ \frac{16\pi m}{r^2}\int_0^r\varphi_0h\,ds\\ \frac{16\pi e}{r^2}\int_0^r\varphi_0h\,ds\end{pmatrix}.
\end{aligned}$$

We observe that, thanks to the nondegeneracy of the solution of Choquard's equation, $D_{\varphi,\chi,\tau,\zeta}(0,\phi_0)$ is a one-to-one operator. Moreover, it can be written as a sum of an isomorphism and a compact operator. It is thus an isomorphism. Finally, the application of the implicit function theorem yields the following result, which is equivalent to Theorem 1.1.

Theorem 2.1. *Suppose $e^2-m^2<0$ and let ϕ_0 be the ground state solution of (12), then there exists $\delta>0$ and a function $\eta\in\mathcal{C}((0,\delta),X_\varphi\times X_\chi\times X_\tau\times X_\zeta)$ such that $\eta(0)=\phi_0$ and $D(\varepsilon,\eta(\varepsilon))=0$ for $0\le\varepsilon<\delta$.*

ACKNOWLEDGEMENTS

The author would like to thank professor Eric Séré and professor Bernhard Ruf for helpful discussions.

REFERENCES

[1] H. Ounaies, *Perturbation method for a class of non linear Dirac equations.* Differential Integral Equations 13 (2000), no. 4-6, 707–720.

[2] E. Lenzmann, *Uniqueness of ground states for pseudo-relativistic Hartree equations.* Preprint (2008).

[3] E. H. Lieb, *Existence and Uniqueness of the Minimizing Solution of Choquard's Nonlinear Equation.* Studies in Applied Mathematics 57 (1977), 93–105.

[4] P.L. Lions, *The Choquard equation and related questions.* Nonlinear Analysis. Theory, Methods and Applications 4 (1980), no. 6, 1063–1073.

[5] S. Rota Nodari, *Perturbation Method for Particle-like Solutions of the Einstein-Dirac Equations.* Ann. Henri Poincaré (2009), DOI 10.1007/s00023-009-0015-x.

[6] F. Finster, J. Smoller, S.T. Yau, *Particlelike solutions of the Einstein-Dirac equations.* Physical Review. D. Particles and Fields. Third Series 59 (1999).

[7] F. Finster, J. Smoller, S.T. Yau, *Particle-like solutions of the Einstein-Dirac-Maxwell equations.* Phys. Lett. A. 259 (1999), no. 6, 431–436.

E-mail address: `rotanodari@ceremade.dauphine.fr`

CEREMADE (UMR CNRS 7534), UNIVERSITÉ PARIS-DAUPHINE, PLACE DU MARÉCHAL DE LATTRE DE TASSIGNY, 75775 PARIS CEDEX 16, FRANCE

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI MILANO, VIA SALDINI 50, 20133 MILAN, ITALY